

~~divided into 3 parts to correspond to its segments, more or less widely margined, translucent or foliaceous, and without teeth.~~

~~Genus *Paracryptocerus* Emery,
 subgenus *Cyathomyrmex* Creighton~~

~~*Cyathocephalus* Emery, 1915, Bul. Soc. Ent. de France,
 p. 192. Preoccupied by Kessler, 1868.~~

~~Type: *Cryptocerus pallens* Klug. By original designation.~~

~~*Cyathomyrmex* Creighton, 1933, Psyche 40: 98. New name.~~

W L Brown Jr C I

STRUMIGENYS VENATRIX WESSON AND WESSON SYNONYMOUS WITH *S. TALPA* WEBER.—In the course of his studies of dacetine ants, Mr. William L. Brown, Jr. secured a loan of the type of *S. talpa* Weber (1934, Psyche, 41: 63-65, fig. 1) from the collections of the Illinois Natural History Survey. This specimen he very kindly placed at my disposal, since I had not seen it during earlier studies on *Strumigenys* in Ohio (Wesson and Wesson, 1939, Psyche, 46: 91-112, Pl. 3). The type of *talpa* proves to be indistinguishable from paratypes of *S. venatrix* which I had described from southern Ohio, and the latter name should be dropped.

According to Brown's recent revision of the dacetine genera, *S. talpa* should be transferred from the genus *Strumigenys* Fred. Smith to the genus *Smithistruma* Brown (1948, Trans. Amer. Ent. Soc. 74: 101-129, 2 figs.).

—LAURENCE G. WESSON, JR., Department of Physiology,
 New York University College of Medicine.